5

10

What is claimed:

- 1. An albumin fusion protein comprising a member selected from the group consisting of:
- (a) a Therapeutic protein:X and albumin comprising the amino acid sequence of SEQ ID NO:18;
 - (b) a Therapeutic protein:X and a fragment or a variant of the amino acid sequence of SEQ ID NO:18, wherein said fragment or variant has albumin activity;
 - (c) a Therapeutic protein:X and a fragment or a variant of the amino acid sequence of SEQ ID NO:18, wherein said fragment or variant has albumin activity, and further wherein said albumin activity is the ability to prolong the shelf life of the Therapeutic protein:X compared to the shelf-life of the Therapeutic protein:X in an unfused state;
 - (d) a Therapeutic protein:X and a fragment or a variant of the amino acid sequence of SEQ ID NO:18, wherein said fragment or variant has albumin activity, and further wherein the fragment or variant comprises the amino acid sequence of amino acids 1-387 of SEQ ID NO:18;
 - (e) a fragment or variant of a Therapeutic protein:X and albumin comprising the amino acid sequence of SEQ ID NO:18, wherein said fragment or variant has a biological activity of the Therapeutic protein:X;
 - (f) a Therapeutic protein:X, or fragment or variant thereof, and albumin, or fragment or variant thereof, of (a) to (e), wherein the Therapeutic protein:X, or fragment or variant thereof, is fused to the N-terminus of albumin, or the N-terminus of the fragment or variant of albumin;
- (g) a Therapeutic protein:X, or fragment or variant thereof, and albumin, or fragment or variant thereof, of (a) to (e), wherein the Therapeutic protein:X, or fragment or variant thereof, is fused to the C-terminus of albumin, or the C-terminus of the fragment or variant of albumin;
 - (h) a Therapeutic protein:X, or fragment or variant thereof, and albumin,

10

or fragment or variant thereof, of (a) to (e), wherein the Therapeutic protein:X, or fragment or variant thereof, is fused to the N- terminus and C-terminus of albumin, or the N-terminus and the C-terminus of the fragment or variant of albumin;

- (i) a Therapeutic protein:X, or fragment or variant thereof, and albumin, or fragment or variant thereof, of (a) to (e), which comprises a first Therapeutic protein:X, or fragment or variant thereof, and a second Therapeutic protein:X, or fragment or variant thereof, wherein said first Therapeutic protein:X, or fragment or variant thereof, is different from said second Therapeutic protein:X, or fragment or variant thereof;
- (j) a Therapeutic protein:X, or fragment or variant thereof, and albumin, or fragment or variant thereof, of (a) to (i), wherein the Therapeutic protein:X, or fragment or variant thereof, is separated from the albumin or the fragment or variant of albumin by a linker; and
- (k) a Therapeutic protein:X, or fragment or variant thereof, and albumin, or fragment or variant thereof, of (a) to (j), wherein the albumin fusion protein has the following formula:

R1-L-R2; R2-L-R1; or R1-L-R2-L-R1,

and further wherein R1 is Therapeutic protein:X, or fragment or variant thereof, L is a peptide linker, and R2 is albumin comprising the amino acid sequence of SEQ ID NO:18 or a fragment or variant of albumin.

- 2. The albumin fusion protein of claim 1, wherein the shelf-life of the albumin fusion protein is greater than the shelf-life of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.
- 3. The albumin fusion protein of claim 1, wherein the in vitro biological activity of the Therapeutic protein:X, or fragment or variant thereof, fused to albumin, or fragment or variant thereof, is greater than the in vitro biological activity of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.

5

10

- 4. The albumin fusion protein of claim 1, wherein the in vivo biological activity of the Therapeutic protein:X, or fragment or variant thereof, fused to albumin, or fragment or variant thereof, is greater than the in vivo biological activity of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.
- 5. An albumin fusion protein comprising a Therapeutic protein:X, or fragment or variant thereof, inserted into an albumin, or fragment or variant thereof, comprising the amino acid sequence of SEQ ID NO:18 or fragment or variant thereof.
- 6. An albumin fusion protein comprising a Therapeutic protein:X, or fragment or variant thereof, inserted into an albumin, or fragment or variant thereof, comprising an amino acid sequence selected from the group consisting of:
 - (a) amino acids 54 to 61 of SEQ ID NO:18;
 - (b) amino acids 76 to 89 of SEQ ID NO:18;
 - (c) amino acids 92 to 100 of SEQ ID NO:18;
 - (d) amino acids 170 to 176 of SEQ ID NO:18;
 - (e) amino acids 247 to 252 of SEQ ID NO:18;
 - (f) amino acids 266 to 277 of SEQ ID NO:18;
 - (g) amino acids 280 to 288 of SEQ ID NO:18;
 - (h) amino acids 362 to 368 of SEQ ID NO:18;
 - (i) amino acids 439 to 447 of SEQ ID NO:18;
 - (j) amino acids 462 to 475 of SEQ ID NO:18;
 - (k) amino acids 478 to 486 of SEQ ID NO:18; and
 - (1) amino acids 560 to 566 of SEQ ID NO:18.
- 7. The albumin fusion protein of claim 5, wherein said albumin fusion protein comprises a portion of albumin sufficient to prolong the shelf-life of the Therapeutic

5

10

protein:X, or fragment or variant thereof, as compared to the shelf-life of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.

- 8. The albumin fusion protein of claim 6, wherein said albumin fusion protein comprises a portion of albumin sufficient to prolong the shelf-life of the Therapeutic protein:X, or fragment or variant thereof, as compared to the shelf-life of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.
- 9. The albumin fusion protein of claim 5, wherein said albumin fusion protein comprises a portion of albumin sufficient to prolong the in vitro biological activity of the Therapeutic protein:X, or fragment or variant thereof, fused to albumin as compared to the in vitro biological activity of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.
- 10. The albumin fusion protein of claim 6, wherein said albumin fusion protein comprises a portion of albumin sufficient to prolong the in vitro biological activity of the Therapeutic protein:X, or fragment or variant thereof, fused to albumin as compared to the in vitro biological activity of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.
- 11. The albumin fusion protein of claim 5 wherein said albumin fusion protein comprises a portion of albumin sufficient to prolong the in vivo biological activity of the Therapeutic protein:X, or fragment or variant thereof, fused to albumin compared to the in vivo biological activity of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.
- 12. The albumin fusion protein of claim 6 wherein said albumin fusion protein comprises a portion of albumin sufficient to prolong the in vivo biological activity of the

10

Therapeutic protein:X, or fragment or variant thereof, fused to albumin compared to the in vivo biological activity of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.

- 5 13. The albumin fusion protein of any one of claims 1-12, which is non-glycosylated.
 - 14. The albumin fusion protein of any one of claims 1-12, which is expressed in yeast.
 - 15. The albumin fusion protein of claim 14, wherein the yeast is glycosylation deficient.
 - 16. The albumin fusion protein of claim 14 wherein the yeast is glycosylation and protease deficient.
 - 17. The albumin fusion protein of any one of claims 1-12, which is expressed by a mammalian cell.
 - 18. The albumin fusion protein of any one of claims 1-12, wherein the albumin fusion protein is expressed by a mammalian cell in culture.
 - 19. The albumin fusion protein of any one of claims 1-12, wherein the albumin fusion protein further comprises a secretion leader sequence.
 - 20. A composition comprising the albumin fusion protein of any one of claims 1-12 and a pharmaceutically acceptable carrier.

- 21. A kit comprising the composition of claim 20.
- 22. A method of treating a disease or disorder in a patient, comprising the step of administering the albumin fusion protein of any one of claims 1-12.

- 23. The method of claim 22, wherein the disease or disorder comprises indication:Y.
- 24. A method of treating a patient with a disease or disorder that is modulated by

 Therapeutic protein:X, or fragment or variant thereof, comprising the step of administering
 an effective amount of the albumin fusion protein of any one of claims 1-12.
 - 25. The method of claim 24, wherein the disease or disorder is indication: Y.
 - 26. A method of extending the shelf life of Therapeutic protein:X, or fragment or variant thereof, comprising the step of fusing the Therapeutic protein:X, or fragment or variant thereof, to albumin, or fragment or variant thereof, sufficient to extend the shelf-life of the Therapeutic protein:X, or fragment or variant thereof, compared to the shelf-life of the Therapeutic protein:X, or fragment or variant thereof, in an unfused state.
 - 27. A nucleic acid molecule comprising a polynucleotide sequence encoding the albumin fusion protein of any one of claims 1-12.
 - 28. A vector comprising the nucleic acid molecule of claim 27.

25

29. A host cell comprising the nucleic acid molecule of claim 28.